



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

specimens of *B. l. lineatus* from Connecticut, having the chest and breast uniform bright cinnamon rufous and the abdomen, tibiae and lower tail-coverts bright buff heavily barred with cinnamon rufous. They are larger than *B. l. alleni* from Florida and have the head and neck not grayish but even more rufous than *lineatus*.

Six young birds collected at the same time differ from the description of young *B. l. elegans* by having the pale spaces on the outer webs of the primaries as large as in *B. l. lineatus*. From the latter they differ by having the lower parts, especially the tibiae, more buffy and the dark markings larger — sagittate or cuneiform instead of oval — and numerous even on the tibiae, which are slightly if at all spotted in *B. l. lineatus*. Young *B. l. alleni* is smaller and has less buff in the plumage, and the dark markings below are even heavier than in the Texas race.

GENERAL NOTES.

Holboëll's Grebe in Connecticut.— An unusual flight of *Colymbus holboëlli* was noticed here during the month of February, 1912. A specimen was picked up alive in a mowing lot, perhaps two miles from the Connecticut river, on the 9th, the ground at the time being covered with snow and the thermometer near the zero point. The bird could not, apparently, arise from the ground. From the 12th to the 15th, inclusive, nine of these grebes were captured alive on the ice in the Connecticut river. Some of them in trying to escape simply moved along the ice in a rapid manner using their feet for power, but making no attempt to fly. Two of them, however, arose from the ice and flew at a height of from eight inches to two feet for a short distance and then dropped down. Between the dates referred to the river was entirely covered with ice, there being, so far as could be seen, no open water where the birds could obtain food.—JNO. H. SAGE, *Portland, Conn.*

The Forked-tailed Gull (*Xema furcatum*).— Recently in looking over some notes taken at sea a number of years ago (1885) and which had been forgotten I came across the following in relation to the Forked-tailed Gull.

In making a passage from Callao, Peru, to Acapulco, Mexico, we passed in sight of Chatham, one of the Galapagos Islands. When three to four hundred miles distant from the island — both when approaching and when